

ACIP-Recommended Immunizations for Adults

The Advisory Committee on Immunization Practices (ACIP) comprises medical and public health experts who develop recommendations on the use of vaccines in the civilian population of the United States. The recommendations stand as public health guidance for safe use of vaccines and related biological products.

These immunizations may be fully covered with no cost to you. Check your coverage with your claims administrator and talk with your healthcare provider about which immunizations are right for you.

Legend:

Recommended vaccination for adults who meet age requirement, lack documentation of vaccination, or lack evidence of past infection	Recommended vaccination for adults with an additional risk factor or another indication	Recommended vaccination based on shared clinical decision-making	No recommendation/Not applicable
--	---	--	----------------------------------

Vaccine	19-26 years	27-49 years	50-64 years	65+ years
Influenza (IIV4, RIV4)	1 dose annually			
Tetanus, Diphtheria, Pertussis (Tdap, Td)	1 dose Tdap, then Td or Tdap booster every 10 years			
	1 dose Tdap each pregnancy; 1 dose Td/Tdap for wound management			
Measles, Mumps, Rubella (MMR)	1 or 2 doses depending on indication (if born in 1957 or later)			
Varicella (VAR)	2 doses (if born in 1980 or later)		2 doses (if born before 1980)	
Zoster Recombinant (RZV)	2 doses for immunocompromising conditions		2 doses	
Human Papillomavirus (HPV)	2 or 3 doses depending on age at initial vaccination or condition	27 through 45 years		
Pneumococcal (PCV15, PCV20, PPSV23)	1 dose PCV15 followed by PPSV23 OR 1 dose PCV20			1 dose PCV15 followed by PPSV23 OR 1 dose PCV20
Hepatitis A (HepA)	2 or 3 doses depending on vaccine			
Hepatitis B (HepB)	2, 3, or 4 doses depending on vaccine or condition (19 through 59 years)			60+ years
Meningococcal A, C, W, Y (MenACWY)	1 or 2 doses depending on indication			
Meningococcal B (MenB)	19 through 23 years	2 or 3 doses depending on vaccine and indication (24+ years)		
Haemophilus Influenzae Type B (Hib)	1 or 3 doses depending on indication			
COVID-19	CDC recommends everyone 5+ years get their primary series of COVID-19 vaccine, and everyone 12+ years also receive a booster shot			

Source: Centers for Disease Control and Prevention ([cdc.gov](https://www.cdc.gov))